


DOING PHYSICS WITH MATLAB

QUANTUM MECHANIC



THEORY OF ALPHA DECAY:

ANALYTICAL and NUMERICAL MODELLING

Ian Cooper School of Physics University of Sydney ian.cooper@sydney.edu.au	Badr Eddine El Fatehy Chfira University of Murcia (UMU, Spain) https://www.um.es/ badreddine.el@um.es
---	---

DOWNLOAD DIRECTORY FOR MATLAB SCRIPTS

qpAlphaDecayEAN.m

The Script can be used to compare Experimental, Analytical and Numerical values for the half-life values as a function of the kinetic energy of the escaped alpha particles for the isotopes of polonium and uranium. The Experimental values of the kinetic energies T and half-lives h are stored in arrays (polonium: kinetic energy $T1_E$; half-life $h1_E$ and uranium: kinetic energy $T2_E$; $h2_E$). The Geiger-Nuttall law is used to calculate the half-life values (Analytical values h_A). A [finite difference method](#) is used to solve the Schrodinger Equation for a nuclear potential function (Numerical values h_N). The Script can easily be changed to model the alpha decay of other elements.

http://www.physics.usyd.edu.au/teach_res/mp/mphome.htm

se_fdm_barrier.m

The wavefunction for a rectangular barrier is computed by solving the Schrodinger Equation by a [finite difference method](#). The code is a modification of the mscript `se_fdm.m`. The height of the barrier is set at 100 eV. The total energy of the system is entered via the Command Window ($E < 100$ eV).

qpAlphaDecayN.m

This Script is used to compute the half-life of an unstable nucleus which emits an alpha particle. The radioactive isotope is specified by its atomic number Z and its mass number A . The half-life $t_{1/2}$ is computed from a given value of the kinetic energy T of the emitted alpha particle. The probability P of the alpha particle tunneling through the potential barrier is found by solving the Schrodinger Equation using a finite difference method.

Input variables:

Atomic number Z ; mass number A ; escape kinetic of alpha particle T ; radius factor Rf (~ 1 to ~ 1.5 . Value can be adjusted to get better agreement between the experimental value of the half-life and the computed value); angular momentum quantum number LA ; switch to choose a square well or the Woods-Saxon nucleus potential `flagWS`; diffuseness parameter a (~ 0.25 to ~ 1); well depth $U0$ ($U0 < 0$); maximum radial coordinate `xMax`; and number of grid points for calculations N ($N = 999999$, need to use a very large number).

Computations:

The Schrodinger equation is solved using a finite difference method to find the wavefunction `psi` and transmission probability P of tunneling through the potential barrier for the selected potential energy function. The results of the computations are displayed in the Command Window as tables and in a graphical output.

http://www.physics.usyd.edu.au/teach_res/mp/mphome.htm

qpAlphaDEcay3D.m

Computation of the half-life for the nuclear potential energy function which includes contributions from Woods-Saxon nuclear potential, centrifugal potential (angular momentum) and Coulomb potential for the repulsion of the alpha particle. The execution of the Script calculates the half-lives for a set of diffuseness parameters and angular momentum values.

qpAlphaDecayWKB.m

Computation of the half-life using the WKB approximation. The WKB approximation is not discussed in the notes.

qpAlphaDecayP.m

The Script uses a finite difference method to calculate the half-life as a function of the alpha particle kinetic energy. The function **polyfit** is used to fit a curve to the data.

ALPHA DECAY


A radioactive substance becomes more stable by: **alpha decay** (helium nucleus ${}^4\text{He}_2$) or **beta decay** (electron e^- or positron e^+) or **gamma decay** (high energy photon). If a radioactive substance at time $t = 0$ contains N_0 radioactive nuclei, then the number of nuclei N remaining after a time interval t is

$$(1) \quad N = N_0 e^{-\gamma t}$$

where γ is the **decay constant** which is the probability per unit time interval that a nucleus will decay. The **half-life** $t_{1/2}$ is defined as the time required for half of a given number of nuclei to decay. The decay constant and half-life are related by

$$(2) \quad t_{1/2} = \frac{\ln(2)}{\gamma}$$

In **alpha decay**, a nucleus emits a helium nucleus ${}^4\text{He}_2$ which consists of 2 protons and 2 neutrons. Therefore, the atomic number Z decreases by 2 and the mass number A decreases by 4. If X is the parent nucleus and Y is the daughter nucleus then the decay can be written as



The attractive forces binding the nucleons together within the nucleus are of short range and the total binding force is approximately proportional to its mass number A . The repulsive electrostatic force between the protons is of an unlimited range and the disruptive force in a nucleus is approximately proportional to the square of the atomic number. Nuclei with $A > 210$ are so large that the short range strong nuclear force barely balances the mutual repulsion of their protons. Alpha decay occurs in such nuclei as a means of reducing their size and increasing the stability of the resulting nuclei after the decay has occurred. A simple model for the potential energy $U(r)$ of the alpha particle as a function of distance r from the centre of the heavy nucleus is shown in figure 1.

The potential well is modelled as a three dimensional deep attractive well associated with the strong nuclear force and beyond the range of this force an electrostatic repulsion due to the Coulomb force between the daughter nucleus and the alpha particle as given by

$$(3) \quad U(r) = \frac{2(Z-2)e^2}{4\pi\epsilon_0 r}$$

where r is the distance from the centre of the nucleus, Z is the atomic number of the parent nucleus and $(Z-2)$ is the atomic number of the daughter nucleus.




Fig. 1. The potential energy U of an alpha particle and nucleus as a function of the distance r from the centre of the nucleus. Total energy E , potential energy U , kinetic energy K and kinetic energy of escaped alpha particle is $T \equiv E_\alpha$. R_0 is the width of the potential well and is equal to radius of the alpha particle plus the radius of the daughter nucleus. R_1 is the radius at which the alpha particle escapes from the parent nucleus ($K = 0$).

From a classical point of view, an alpha particle does not have sufficient energy to escape from the potential well. The barrier height is typically in the order of 30 MeV and the decayed alpha particles have energies only in the range 4 to 9 MeV. This range of energies from 4 to 9 MeV is very small, but the half-lives of the radioactive alpha emitters range from $\sim 10^{-7}$ s to $\sim 10^{18}$ s, while the energy changes by a factor of 2, the change in the half-lives is about **25 orders of magnitude !!!**. Our goal is explain, how an alpha particle can escape from a nucleus and find a relationship between the energy of the alpha particle and its half-life.

We will do this by (1) Analytical means and (2) Numerically by solving the Schrodinger equation using a finite difference method for an alpha particle in a potential well as shown in figure (1).

Classical arguments fail to account for alpha decay, but quantum mechanics provides a straight forward explanation based upon the concept of **tunnelling** where a particle can be found in a classically forbidden region. The theory of alpha decay was developed in 1928 by Gamow, Gurney and Condon and provided confirmation of the power of quantum mechanics. We will develop simplified models to account for alpha decay and compare our predictions with those of experiments.

ANALYTICAL MODEL

Reference: *Concepts of Modern Physics*, Arthur Beiser

Assume that an alpha particle exists as a particle in constant motion with velocity v_α contained inside the heavy nucleus by a surrounding potential barrier. There is a small probability that the alpha particle can escape each time it encounters the barrier. The decay constant γ gives the decay probability per unit time of an escape

$$(4) \quad \gamma = f P$$

where f is the number of times per second an alpha particle strikes the potential barrier at R_0 and P is the probability that the particle passes through the barrier. The probability can be determined from the wavefunction. Inside the potential well ($0 \leq r \leq R_0$) the wavefunction has an oscillatory nature and its amplitude is A_{in} . In the barrier region ($R_0 < r \leq R_1$) the wavefunction decreases exponentially. When the alpha particle has escaped ($r > R_1$) the wavefunction is approximately sinusoidal with an amplitude A_{out} , therefore, the probability P that the alpha particle strikes the barrier and escapes is

$$(5) \quad P = \left(\frac{A_{out}}{A_{in}} \right)^2$$

An example of tunnelling through a barrier is shown in figure 2. Classically the particle does not have sufficient energy to pass through the barrier and upon striking the barrier the particle would be reflected.




Fig. 2. The wavefunction for a barrier of finite width. The particle is incident upon the barrier from the left. There is a small probability that the particle is located to the right of the barrier. [se_fdm_barrier.m](#)

Assume that the alpha particle moves back and forth in the nucleus of radius R_0 . We can estimate the radius R_0 assuming that it is equal to the radius of the alpha particle plus the daughter nucleus (figure 1)

$$(6) \quad R_0 = R_F (4^{1/3} + (A-4)^{1/3}) \text{ fm}$$

where A is the mass number of the parent nucleus and R_F is a radius factor. The nuclear radius R_0 is not a well-defined quantity. It is often found by fitting experimental data to a theoretical fit when $\ln(t_{1/2})$ is plotted against $1/\sqrt{T}$.

Then the frequency f of striking the barrier is

$$(7) \quad f = \frac{v_{in}}{2R_0}$$

where v_{in} is the alpha particle velocity inside the parent nucleus

$$(8) \quad v_{in} = \sqrt{\frac{T - U_0}{2m_R}}$$

The alpha particle goes one way while the daughter nucleus recoil in the opposite direction. So, it may be better to use the reduced mass m_R rather than the mass of the alpha particle m_α where

$$m_\alpha = m_R (A - 4) / A$$

For the barrier region, $U > E$ and $K < 0$, classical physics predicts a transmission probability $P = 0$. However, in the quantum picture, the particle behaves as a wave and there is a non-zero probability of the particle escaping from the potential well, $P > 0$. The particle escapes the barrier at $U = E$ at the position R_1 where

$$(9) \quad R_1 = \frac{2(Z - 2)e^2}{4\pi\epsilon_0 E_\alpha} \quad U(R_1) = E_\alpha = T$$

After much “hard work” and assuming that $R_0 \ll R_1$ it can be shown that the probability of an alpha particle escaping our potential well is

(10)

$$P = \exp \left[\left(\frac{4e}{\hbar} \right) \left(\frac{m_R}{\pi \epsilon_0} \right)^{1/2} R_0^{1/2} (Z-2)^{1/2} - \left(\frac{e^2}{\hbar \epsilon_0} \right) \left(\frac{m_R}{2} \right)^{1/2} (Z-2) T^{-1/2} \right]$$

We are now able to estimate the half-life of an alpha emitted from an unstable nucleus given the atomic number Z and the mass number A of the parent nucleus and the mass m_R and kinetic energy T of the escaped alpha particle.

The half-life $t_{1/2}$ can be computed from the known parameters of Z , A , T , v_{in} and R_0 .

1: Calculate R_0 from Eq. (6) $R_0 = R_f \left(4^{1/3} + (A-4)^{1/3} \right) \times 10^{-15} \text{ m}$

2: Calculate v_α from Eq. (8) $v_{in} = \sqrt{\frac{T - U_0}{2m_R}}$

3: Calculate f from Eq. (7) $f = \frac{v_\alpha}{2R_0}$

4: Calculate P from Eq. (10)

$$P = \exp \left[\left(\frac{4e}{\hbar} \right) \left(\frac{m_R}{\pi \epsilon_0} \right)^{1/2} R_0^{1/2} (Z-2)^{1/2} - \left(\frac{e^2}{\hbar \epsilon_0} \right) \left(\frac{m_R}{2} \right)^{1/2} (Z-2) T^{-1/2} \right]$$

http://www.physics.usyd.edu.au/teach_res/mp/mphome.htm

5: Calculate γ from Eq. (4) $\gamma = f P$

6: Calculate $t_{1/2}$ from Eq. (2) $t_{1/2} = \frac{\ln(2)}{\gamma}$

We can rewrite the expression given by equation 6 for the half-life as

$$P = \exp\left(K_1 - K_2 T^{-1/2}\right)$$

$$K_1 = \left(\frac{4e}{\hbar}\right) \left(\frac{m_R}{\pi \epsilon_0}\right)^{1/2} R_0^{1/2} (Z-2)^{1/2}$$

$$K_2 = \left(\frac{e^2}{\hbar \epsilon_0}\right) \left(\frac{m_R}{2}\right)^{1/2} (Z-2)$$

$$t_{1/2} = \frac{\ln(2)}{\gamma}$$

$$\ln(t_{1/2}) = \ln\left(\frac{\ln(2)}{\gamma}\right) = \ln(2) - \ln(\gamma) = \ln(2) - \ln(f P)$$

$$\ln(t_{1/2}) = \ln(2) - \ln(f P) = \ln\left(\frac{2}{f}\right) - K_1 + K_2 T^{-1/2}$$

$$(11) \quad \ln(t_{1/2}) = \left(\ln\left(\frac{2}{f}\right) - K_1\right) + K_2 T^{-1/2}$$

Equation (11) is approximately a straight line when $\ln(t_{1/2})$ [y-axis] is plotted against $1/T^{-1/2}$ [x-axis].

$$\ln(t_{1/2}) = b + m T^{-1/2} \quad (y = b + m x)$$

This expression is known as the **Geiger-Nuttall rule**.

http://www.physics.usyd.edu.au/teach_res/mp/mphome.htm

NUMERICAL ANALYSIS

We can compute the half-life $t_{1/2}$ for the radioactive decay of a nucleus which emits an alpha particle by solving the Schrodinger Equation by a [finite difference method](#) to find the wavefunction as a function of the radial coordinate. The Script **qpAlphaDecayN.m** can be used to solve the Schrodinger Equation and compute the half-life.

We need to solve the Schrodinger Equation for the a potential energy function representing the system of the parent nucleus and the alpha particle (figure 1) to find the amplitude A_{in} of the wavefunction inside the potential well ($r < R_0$) and the amplitude A_{out} of the wavefunction outside the well ($r > R_1$). Once, A_{in} and A_{out} have been computed, then it east to calculate the half-life $t_{1/2}$ as was describe in the Analytical method.

The **time-independent Schrodinger Equation in one dimension** can be expressed as (x is used as the radial coordinate and not r)

$$(13) \quad -\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + U(x) \psi(x) = E \psi(x)$$

For nuclear systems it is more convenient to measure lengths in fm (femometre) and energies in MeV. We can use the scaling factors

length: $L_{se} = 1 \times 10^{-15}$ to convert m into fm

energy: $E_{se} = 1.6 \times 10^{-13}$ to convert J into MeV

So, we can write equation (13) as

http://www.physics.usyd.edu.au/teach_res/mp/mphome.htm

$$(14) \quad \left[\left(\frac{-\hbar^2}{2m} \right) \left(\frac{1}{L_{se}^2 E_{se}} \right) \frac{d^2}{dx^2} + U(x) \right] \psi(x) = E \psi(x)$$

$$\left[C_{se} \frac{d^2}{dx^2} + U(x) \right] \psi(x) = E \psi(x) \quad \text{where} \quad C_{se} = \left(\frac{-\hbar^2}{2m} \right) \left(\frac{1}{L_{se}^2 E_{se}} \right)$$

where m is the reduced mass.

In the finite difference method, the second derivative of the wavefunction is approximated by

$$(16) \quad \frac{d^2 \psi(x)}{dx^2} = \frac{\psi(x + \Delta x) - 2\psi(x) + \psi(x - \Delta x)}{\Delta x^2}$$

The Schrodinger Equation must be solved backwards starting from a far-away point.

```
% Wavefunction
for n = N-1:-1:2
    SEconst = (T - U(n)) .* dx^2 ./ Cse;
    psi(n-1) = (2 - SEconst) * psi(n) - psi(n+1);
end

psi = psi ./ max(psi);
```

The point at which the potential is equal to the kinetic energy of the alpha particle is found using the Matlab function **solve**

```
% Second turning point for each value of the orbital
angular momentum, LA= 0, 1, 2, 3
% T = U
x1 = max(double(solve(@(z) LA.*(LA+1).*Cse./(z.^2) +
2.*(Z-2).*e.^2./ (4.*pi.*eps0.*Ese.* Lse.* z) ==
T))); % U(R1) = T
```

The amplitudes of the wavefunction inside and outside the potential well are found by using Matlab **find** function from which the half-life is computed. The kinetic energy of the alpha particle can be calculated from the total energy and from the wavelength of the wavefunction using the Matlab function **findpeaks**

```
% CALCULATIONS =====
% Position in arrays
% index for x position when E = U outside well
    index = find(x>x1, 1 );
% index for position outside well
    index1 = find(x>x0, 1 );
% index for position to calc Aout
    index2 = round(0.9*N);
% velocity of alpha particle inside well
    v_in = sqrt(2*(T-U0)*Ese/mR);
% frequency at which particles strike barrier on the RHS of well
    f = v_in / (2*x0*Lse);
% amplitude of wavefunction inside and outside nucleus
    Ain = max(abs(psi(1:index1)));
    Aout = max(abs(psi(index2:N)));
% probability of alpha particle escaping
    P = (Aout / Ain)^2;
% decay constant
    gamma = f * P;
% half-life
    half_life = log(2) / gamma;

% Wavelength --> kinetic energy
    [~, z2] = findpeaks(psi(1:index1));
    wLin = (x(z2(end))-x(z2(1)))/(length(z2)-1);
    [z1, z2] = findpeaks(psi(index2:N));
    wLout = (x(z2(end)+index2)-x(z2(1)+index2))/(length(z2)-1);

    p_in = h/(wLin*1e-15);
    T_in = p_in^2/(2*mR)/(e*1e6);

    p_out = h/(wLout*1e-15);
    T_out = p_out^2/(2*mR)/(e*1e6);

% kinetic energy T - U
    K_in = T - U0;
    K_out = T - U(end);

    Umax = max(U);
```

MATLAB SIMULATIONS

qpAlphaDecayEAN.m

Comparison of the Experimental, Analytical values and Numerical values for the half-lives of polonium and uranium isotopes.

The results are displayed in the Command Window using the Matlab function **table**.

The tables columns are A (mass numbers), T (kinetic energy of escaped alpha particle), experimental half-lives (h_E), analytical half-lives (h_A), ratio of experimental to analytical half-lives (R_EA), numerical half-lives (h_N) and ratio of experimental to numerical half-lives (R_EN). The energies are in MeV and half-lives in s.

Isotopes of polonium

A1	TP	hP_E	hP_A	RP_EA	hP_N	RP_EN
218	6.002	1.86e+02	2.8e+02	0.7	4.5e+02	0.4
217	6.537	1.53e+00	1.2e+00	1.3	3.6e+00	0.4
216	6.778	1.50e-01	1.3e-01	1.1	5.2e-03	28.8
215	7.370	1.80e-03	8.6e-04	2.1	1.8e-02	0.1
214	7.687	1.62e-04	7.6e-05	2.1	1.1e-03	0.1
213	8.376	3.70e-06	6.1e-07	6.1	2.2e-05	0.2
212	8.785	3.00e-07	4.6e-08	6.5	1.1e-06	0.3
210	5.304	1.20e+07	1.7e+06	7.0	6.7e+05	17.9
209	4.883	3.22e+09	6.6e+08	4.9	5.4e+08	5.9
208	5.110	9.15e+07	2.6e+07	3.5	5.0e+07	1.8

Notice the enormous range of half_lives values. For T = 4.883 MeV $t_{1/2} \sim 10^9$ s. However, when T = 8.785 MeV, $t_{1/2} \sim 10^{-7}$ s. For a doubling of the alpha particle energy, the half-life decreases by ~ 16 orders of magnitude.

Isotopes of uranium

A2	TU	hU_E	hU_A	RU_EA	hU_N	RU_EN
238	4.151	1.41e+17	3.1e+17	0.4	1.2e+16	11.7
236	4.445	7.39e+14	1.0e+15	0.7	1.2e+14	6.0
235	4.215	2.22e+16	9.6e+16	0.2	3.0e+17	0.1
234	4.722	7.75e+12	7.4e+12	1.0	3.9e+13	0.2
233	4.729	5.02e+12	6.9e+12	0.7	3.0e+13	0.2
232	5.236	2.23e+09	2.3e+09	1.0	1.3e+10	0.2
230	5.818	1.80e+06	8.9e+05	2.0	8.2e+06	0.2
228	6.410	5.46e+02	9.5e+02	0.6	4.9e+03	0.1
227	6.860	6.60e+01	9.5e+00	7.0	1.3e+02	0.5
226	7.402	2.60e-01	6.4e-02	4.0	8.7e-01	0.3
225	7.875	8.40e-02	1.3e-03	65.4	1.2e-02	6.8
223	8.780	1.80e-05	1.8e-06	10.1	9.7e-05	0.2

Generally, the agreement between the experimental values and the computed values is within an order of magnitude.

The values for the radius factor R_f were adjusted to give a best fit between the experimental values and the computed values.

polonium $R_f = 1.07$ uranium $R_f = 1.15$

There is some doubt over the experimental values quoted. When searching the internet it was difficult to find consistent value. The sources of the experimental data were

http://www.kavelaby.npl.co.uk/atomic_and_nuclear_physics/4_6/4_6_1_part08_080_089.html

http://www.kavelaby.npl.co.uk/atomic_and_nuclear_physics/4_6/4_6_1_part09_090_099.html#Pa

http://www.physics.usyd.edu.au/teach_res/mp/mphome.htm

The results are also displayed in two Figure Windows (figures 3 and 4)




Fig. 3. The straight lines show the predictions of the Geiger-Nuttall rule: $\ln(t_{1/2}) = b + m T^{-1/2}$ ($y = b + m x$). The open circles are the Experimental values, the coloured circles are the Analytical values and the squares are the Numerical values. [qpAlphaDecayEAN.m](#)




Fig. 4. The Geiger-Nuttall relationship for the isotopes of polonium and uranium. The circles are the experimental values.
[qpAlphaDecayEAN.m](#)

How good is the fit? The agreement is really excellent which is quite an achievement considering the sensitive of the results on the transmission probability with the kinetic energy of the ejected alpha particle. Quantum mechanics via the concept of tunnelling can account for the fact that a doubling of the alpha particle energy leads to a change in half-life of 25 orders of magnitude. It is not possible to predict with great accuracy the computed value of a half-life because of many factors such as: the radius of an isotope is not an exact quantity (some uncertainly in both the radius factor R_f and the width of the nuclear well); the shape of the nuclear potential; the energy spectrum of the alpha emitters has a fine structure because an alpha-active nucleus may go over to any number of several different quantized energy levels of its daughter with each having a different energy and the emitted alpha particle may have a non-zero angular momentum.

http://www.physics.usyd.edu.au/teach_res/mp/mphome.htm

qpAlphaDecayN.m

The Script is used to solve the Schrodinger Equation for different nuclear potential energy models to compute the half-life of the specified isotope. The variable **flagWS** is used to select either the square well potential or the Woods-Saxon and you can change the angular momentum **LA** of the emitted alpha particle. The input variables are changed within the Script.

```
% INPUTS =====  
  
% Atomic number of element  
Z = 92;  
% Mass number of element  
A = 236;  
% Kinetic energy of escaped alpha particle [MeV]  
T = 4.445;  
% Nuclear radius factor polonium (1.07) uranium (1.15)  
Rf = 1.15;  
  
% Angular momentum quantum number LA [0, 1, 2, ... ]  
LA = 0;  
  
% Woods Saxon Nuclear Potential flagWS = 1 / square well flagWS =  
0;  
% diffuseness parameter a [0.25 to 1]  
flagWS = 0;  
a = 0.5;  
  
% Depth of potential well [-40 MeV]  
U0 = -40;  
% Max radial coordinate [250 fm]  
xMax = 250;  
% Number of grid points  
N = 1e6;
```

A summary of the computed values is displayed in a table in the Command Window. For example, the isotope of uranium 236:

atomic number $Z = 92$

mass number $A = 236$

kinetic energy (escaped alpha particle) $T = 4.445 \text{ MeV}$

radius factor $R_f = 1.15$

calculations $N = 1 \times 10^6$

diffuseness $a = 0.5$

half-life $= 5.5082 \times 10^{14} \text{ s}$

gamma $= 1.1934 \times 10^{-15} \text{ s}^{-1}$

tunneling probability $P = 4.4583 \times 10^{-37}$

radius potential well $x_0 = 8.8918 \text{ fm}$

max potential energy $U_{\text{max}} = 29.182 \text{ MeV}$

radial position when $K = 0$ $x_1 = 58.378 \text{ fm}$

velocity inside well $v_{\text{in}} = 4.6662 \times 10^7 \text{ m.s}^{-1}$

collision frequency $f = 2.6239 \times 10^{21} \text{ s}^{-1}$

From wavelength estimates

$$T_{\text{in}} = 4.4387 \text{ MeV} \quad T_{\text{out}} = 3.3597 \text{ MeV}$$

From $K = T - U$ estimates

$$K_{\text{in}} = 4.4445 \text{ MeV} \quad K_{\text{out}} = 3.407 \text{ MeV}$$




Fig. 5. Graphical output for the input parameters of the element uranium: $A = 236$ and $Z = 92$ and $T = 4.45$ MeV. The top plot shows the potential energy function (red) and the kinetic energy of the escaped alpha particle (blue). The lower plot shows the wavefunction. Note the enormous magnification factor for the wavefunction inside and outside the potential well, indicating the extremely small probability of the alpha particle tunnelling through the barrier. `qpAlphaDecayN.m`

How do slight changes in the numerical model with the square potential well and the Coulomb potential as shown in figure1 affect the half-life value?

We will consider the results for the isotope of polonium $^{236}\text{U}_{92}$ with the default values and change only a single variable in any one simulation.

Default values

Z = 92; A = 2236; **$t_{1/2}(\text{experiment}) = 7.39 \times 10^{14} \text{ s}$**
T = 4.445; Rf = 1.15, LA = 0; flagWS = 0; U0 = -40; xMax = 250
N = 1×10^6
 $t_{1/2} = 5.8 \times 10^{14} \text{ s}$

$$N = 1 \times 10^4 \Rightarrow t_{1/2} = 5.2 \times 10^{14} \text{ s}$$

$$Rf = 1.25 \Rightarrow t_{1/2} = 2.5 \times 10^{14} \text{ s} \quad Rf = 1.07 \Rightarrow t_{1/2} = 8.1 \times 10^{14} \text{ s}$$

Decreasing the value of Rf increases the estimate of the half-life value.

$$U0 = -30 \Rightarrow t_{1/2} = 7.3 \times 10^{14} \text{ s}$$

$$x\text{Max} = 125 \Rightarrow t_{1/2} = 2.4 \times 10^{13} \text{ s}$$

$$T = 4.8 \Rightarrow t_{1/2} = 7.70 \times 10^{12} \text{ s} \quad \text{half-life very sensitive to T value}$$

NUCLEAR POTENTIAL ENERGY FUNCTION

The alpha decay process involves an interaction between the wavefunction and the nuclear potential energy function. The nuclear potential energy function $U(r)$ can be approximated by the summation of the functions for the potential well $U_N(r)$, the Coulomb repulsion $U_C(r)$ and a centrifugal term $U_L(r)$.

$$U(r) = U_N(r) + U_C(r) + U_L(r)$$

where $U_L(r) = \frac{\hbar l(l+1)}{2m_R r^2}$

l is the angular momentum quantum number ($l = 0, 1, 2, \dots$)

$$U_C(r) = \frac{2(Z-2)e^2}{4\pi\epsilon_0 r} \quad r > R$$

In figure 1, the nuclear potential was a simple square well function. A more complex model is the shape of the nuclear potential energy function is given by the Woods-Saxon curve


$$U_N(r) = \frac{1}{1 + \exp\left(\frac{r - R_0}{a}\right)} \quad \text{diffuseness parameter } a$$

The theory in the Analytical section presented above neglects the effects of angular momentum in that it assumes the alpha particle carries off no orbital angular momentum ($l = 0$). If alpha decay takes place to or from an excited state, some angular momentum may be carried off by the particle with a resulting change in the half-life. For $l > 0$, the alpha particle must tunnel through a wider well and higher barrier.

http://www.physics.usyd.edu.au/teach_res/mp/mphome.htm

The following plots created (`qpAlphaDecayN.m`) show the changes in the nuclear potential energy and half-life for $l > 0$ and using the Woods-Saxon potential (`flagWS = 1`) for the uranium isotope $Z = 92$, $A = 236$, $T = 4.445$ MeV.

Square well $l = 0$




$t_{1/2} = 5.8 \times 10^{14}$ s $x_0 = 8.89$ fm $x_1 = 58.38$ $U_{\max} = 29.18$ MeV

Square well $l = 5$

$t_{1/2} = 4.1 \times 10^{16}$ s $x_0 = 8.89$ fm $x_1 = 58.99$ $U_{\max} = 31.12$ MeV

http://www.physics.usyd.edu.au/teach_res/mp/mphome.htm

Woods-Saxon $a = 1$ $l = 0$



$t_{1/2} = 3.7 \times 10^{13}$ s $x_0 = 8.89$ fm $x_1 = 58.84$ $U_{\text{max}} = 19.92$ MeV

Woods-Saxon $a = 1$ $l = 5$

$t_{1/2} = 2.9 \times 10^{15}$ s $x_0 = 8.89$ fm $x_1 = 58.99$ $U_{\text{max}} = 21.07$ MeV


http://www.physics.usyd.edu.au/teach_res/mp/mphome.htm

You can use the Script **qpAlphaDecay3D.m** to compute the half-lives using the Wood-Saxon with many different a and l values in the one execution of the Script.

Run the Script **qpAlphaDecayWKB.m** to compute the half-life using the WKB approximation. Typical parameters:

$$E=8.78; A=212; Z=84; U_0=0; L=0 \rightarrow 2.4317e-07 \text{ s.}$$

The experimental value is $3.0e-07 \Rightarrow$ quite good agreement



WKB approximation:

$$\text{Inputs: } Z = 84 \quad A = 212 \quad E_\alpha = 8.78 \quad l = 3 \quad U_0 = 40$$


$$\text{Outputs: } R_0 = 9.01 \quad R_1 = 27.20 \quad b = 18.18 \quad vA = 4.85e7$$

$$f = 2.69e21 \quad P = 6.55e-15 \quad \text{lamba} = 1.76e7 \quad t_{1/2} = 3.93e-8$$

http://www.physics.usyd.edu.au/teach_res/mp/mphome.htm

qpAlphaDecayP.m

The Script uses the **polyfit** function to fit a curve to the data.



REFERENCES

http://ocw.mit.edu/courses/electrical-engineering-and-computer-science/6-728-applied-quantum-and-statistical-physics-fall-2006/study-materials/wkb_tunneling2.pdf

<https://fenix.tecnico.ulisboa.pt/downloadFile/3779576931836/Ex8Serie1/>.

<http://hal.in2p3.fr/in2p3-00637971/document>

https://www.researchgate.net/publication/239535664_a_particle_preformation_in_heavy_nuclei_and_penetration_probability

http://www.physics.usyd.edu.au/teach_res/mp/mphome.htm